

Perioperative Glucocorticoid Coverage

A Reassessment 42 Years After Emergence of a Problem

Michael Salem, M.D.,* Robert E. Tainsh, Jr., M.D., M.P.H.,† Jonathan Bromberg, M.D., Ph.D.,‡
D. Lynn Loriaux, M.D., Ph.D.,§ and Bart Chernow, M.D., F.A.C.P.||

From the Surgical Intensive Care Division, Department of Anesthesiology and Critical Care Medicine, the Johns Hopkins University School of Medicine, Baltimore, Maryland; the Department of Anesthesia, Florida Hospital Medical Center, Orlando, Florida;† the Department of Surgery, Medical University Hospital, Medical University of South Carolina, Charleston, South Carolina;‡ the Endocrine Division, Department of Medicine, Oregon Health Science University, Portland, Oregon;§ and the Department of Medicine, Sinai Hospital of Baltimore, and the Departments of Medicine, Anesthesiology, and Critical Care Medicine, the Johns Hopkins University School of Medicine, Baltimore, Maryland||*

Objective

The authors review the historical basis for the provision of perioperative glucocorticoid coverage, and detail the evolution in the understanding of the role of the hypothalamic-pituitary-adrenal (HPA) axis in response to physical stressors. New recommendations are proposed for glucocorticoid-dependent patients who require anesthesia and surgery.

Summary Background Data

In 1952, a patient developed surgery-associated adrenal insufficiency as a result of preoperative withdrawal from glucocorticoid therapy. That case report, and one other in the ensuing 12 months, prompted the publication of recommendations for perioperative glucocorticoid coverage, which became the standard of care. The understanding of the role of the HPA axis in the stress response has been subsequently refined; however, recommendations for perioperative glucocorticoid coverage have not been altered in parallel.

Methods

Studies were identified beginning with the first reports of the physiologic actions of the adrenal glands (1855) and the description and clinical use of cortisone (1930-1993). Studies were selected for review if they were related to or evaluated the provision of stress-related glucocorticoid administration. All clinical studies were evaluated to determine the basis for the provision of perioperative glucocorticoid coverage and the validity of the data used to justify these conclusions.

Conclusions

Clinical and experimental evidence support the concept that the current amount of perioperative glucocorticoid coverage is excessive and has been based on anecdotal information. New recommendations are proposed which suggest that the amount and duration of glucocorticoid coverage should be determined by: a) the preoperative dose of glucocorticoid taken by the

patient, b) the preoperative duration of glucocorticoid administration, and c) the nature and anticipated duration of surgery.

HISTORICAL PERSPECTIVE

Forty-two years have passed since Fraser and colleagues described a patient who developed perioperative circulatory shock as a consequence of secondary adrenal insufficiency.¹ Although primary adrenal insufficiency had been known since the mid 19th century (Sir Thomas Addison's report of adrenal gland destruction by tuberculosis was published in 1855²), the concept of secondary adrenal insufficiency was not developed until nearly a century later. Cushing is credited for the conceptual association of pituitary and adrenal cortical function.^{3,4} Over the next 20 years, Swingle and colleagues⁵⁻⁹ showed that adrenalectomized dogs were predisposed to hypovolemic and laparotomy-induced circulatory shock; they also showed that this could be prevented by administering glucocorticoids.⁹

Although physiologic responses to adrenal medullary extracts had been evaluated by the beginning of the twentieth century,^{10,11} it was not until 1930 that Swingle and Pfiffner showed that adrenal cortical extracts could maintain life in adrenalectomized animals.^{12,13} Reichstein in Switzerland,^{14,15} and Kendall in the United States^{16,17} led the effort to isolate, characterize, and synthesize cortisone and other adrenal steroids, such as deoxycortisone. Swingle then showed that cortisol prevented the circulatory collapse associated with adrenalectomy.⁸ In 1949, Hench and co-workers^{18,19} described the beneficial effects of cortisone in patients with rheumatoid arthritis (Hench shared the Nobel Prize in medicine with Kendall and Reichstein for this contribution). The era of glucocorticoid therapy for diseases other than primary adrenal insufficiency began with this finding.

In 1952, Fraser and co-workers described the first (reported) patient who developed surgery-associated adrenal insufficiency as a consequence of preoperative withdrawal from glucocorticoid therapy.¹ The following year, Lewis and colleagues²⁰ described a similar patient who died several hours after surgery had been performed to repair a flexion contracture of the right knee. The patient had chronic disabling rheumatoid arthritis, and had been receiving cortisone daily for 5 months before the knee surgery. The cortisone was discontinued the day before surgery. A postmortem examination revealed diffuse atrophy and hemorrhage in the adrenal glands. The case report concluded with a list of recommendations for

perioperative glucocorticoid treatment, which became the standard of therapy for perioperative glucocorticoid coverage.²⁰ Those recommendations, amounting to roughly a fourfold increase in glucocorticoid administration, have been generalized so that the current dose of glucocorticoids is quadrupled routinely. For example, a patient receiving 50 mg of prednisone daily for chronic obstructive pulmonary disease can receive 200 mg of prednisone daily during the perioperative period. This is equivalent to 1000 mg of cortisol each day. For the sake of perspective, it is worth noting that the average cortisol production rate in patients with Cushing's syndrome is 36 mg/day.²¹ This excess of glucocorticoid leads to adverse clinical consequences, such as reduced tissue repair rates, decreased glucose tolerance, and increased susceptibility to infection caused by immune compromise. Knowledge of adrenal cortical responses to physical stressors has been refined over the past 20 years.²²⁻²⁶ As a consequence, perioperative glucocorticoid management can be prescribed in a more rational fashion.

This article considers the following issues: 1) evidence supporting the need for perioperative glucocorticoid coverage; 2) identifying patients needing coverage; 3) and prescribing guidelines for perioperative coverage.

THE NEED FOR PERIOPERATIVE GLUCOCORTICOID COVERAGE

The hormonal responses to surgery,²²⁻²⁵ critical illness, and trauma²⁶⁻²⁹ have been studied extensively over the past 20 years. Plasma adrenocorticotrophic hormone (ACTH)³⁰⁻³¹ and cortisol concentrations,^{22,32-36} urinary-free cortisol,^{37,38} and the concentrations of urinary metabolites of cortisol^{39,40} increase in response to surgery, trauma, and other forms of critical illness.^{26,41} There is considerable inter-individual variation in the hypothalamic-pituitary-adrenal cortical (HPA) response to surgical stress.²³ Some of the variation can be attributed to the actions of anesthetics,^{23,42,43} exogenous and endogenous⁴⁴ opiate analgesics (administered intrathecally,⁴⁵ epidurally,⁴⁶ or parenterally⁴⁷), antihypertensive agents,⁴⁸ age,⁴⁹ sleep,⁵⁰ and infection.⁵¹ All of these factors can alter the HPA response to surgery and other stresses. In addition, the pulsatile secretory patterns of ACTH and cortisol⁵² contribute to the apparent variability in the measured responses to surgical stress.

Perioperative adrenal insufficiency is an excellent model for studying stress-associated adrenal crisis. It is an uncommon complication of surgery. Mohler and colleagues⁵³ noted only one case of perioperative adrenal

Address reprint requests to Bart Chernow, Sinai Hospital of Baltimore, 2401 W. Belvedere Avenue, Baltimore, MD 21215-5271.

Accepted for publication October 11, 1993

Table 1. CLINICAL DATA IN GLUCOCORTICOID TREATED PATIENTS IN WHOM POSTOPERATIVE DEATH OR HYPOTENSION WAS PUTATIVELY SECONDARY TO STRESS-INDUCED ADRENOCORTICAL INSUFFICIENCY*

Authors	No. of Cases	Interval between Surgery and Shock or Death (hrs)	Plasma Cortisol (mmol/L)	Supplementary Glucocorticoids	Comments and Observations on the Diagnosis of Acute Stress-Induced Adrenal Insufficiency
Fraser et al. (1952)	1	3	—	No	Inconclusive, fever, dyspnea, death after transfusion, anaphylactic reaction?
Lewis et al. (1953)	1	6	—	No	Not rejected, no biochemical evidence
Salassa et al. (1953)	1	30	—	Yes	Inconclusive, death
Salassa et al. (1953)	1	16	—	No	Not rejected, no biochemical evidence
Downs and Cooper (1955)	1	14	—	Yes	Inconclusive, perforated ulcer, death
Dragsted et al. (1955)	2	8 and 30	—	No	Inconclusive, sudden death
Kittredge (1955)	1	48	—	Yes	Inconclusive, sudden death
Allanby (1957)	1	20	—	No	Inconclusive, perforated colon, death
Kern (1957)	1	23	—	No	Inconclusive, death
Sloynay and Brooke (1957)	1	13	—	No	Inconclusive, death
Winstone and Brooke (1961)	1	24	—	No	Inconclusive, ulcerative colitis, death
Haller and Kirchoff (1963)	1	36	—	No	Inconclusive, death
Downs and Cooper (1955)	1	2	—	No	Inconclusive, hypotension
Marnegel and Kramer (1955)	1	12	—	No	Inconclusive, hypotension
Nicholas et al. (1955)	1	1	—	Yes	Inconclusive, hypotension
Hayes and Kushlan (1956)	4	Preoperative	—	No	Inconclusive, hypotension
Hayes (1956)	5	Preoperative	—	No	Inconclusive, hypotension
Hayes (1956)	6	Preoperative	—	Yes	Inconclusive, hypotension
Howland et al. (1956)	1	Preoperative	—	No	Inconclusive, hypotension
Richman et al. (1956)	1	—	—	No	Inconclusive, hypotension
De Mowbroy (1957)	1	—	—	Yes	Inconclusive, hypotension
Dundee (1957)	2	2 and preoperative	—	Yes	Inconclusive, hypotension
Nicholas et al. (1957)	3	Postoperative	—	Yes	Inconclusive, hypotension
Sloynay and Brooke (1957)	1	12	—	Yes	Inconclusive, hypotension
Sloynay and Brooke (1957)	3	Postoperative, 6, 24	—	No	Cannot be rejected, hypotension
Gillies (1958)	1	Preoperative	—	Yes	Inconclusive, hypotension
Gillies (1958)	1	Preoperative	—	No	Inconclusive, hypotension
Marks et al. (1959)	1	2	390	No	Not adrenal insufficiency, hypotension
Marks et al. (1959)	1	24	520	Yes	Not adrenal insufficiency, hypotension
Clin Anes Conf (1961)	1	24	—	Yes	Inconclusive, hypotension
Sampson et al. (1961)	1	Preoperative	140	No	Fulfill diagnostic criteria of adrenal insufficiency, hypotension
Sampson et al. (1962)	1	Preoperative	140	No	Inconclusive, hypotension
Salem and Lund (1962)	1	Preoperative	—	No	Inconclusive, hypotension
Salem and Lund (1962)	1	Preoperative	—	No	Inconclusive, hypotension
Kaalund et al. (1966)	3	Preoperative and postoperative	—	Yes	Inconclusive, hypotension
Jansani et al. (1968)	1	Preoperative	170	No	Fulfill the diagnostic criteria
Kehlet and Binder (1973)	1	Preoperative	60	No	Fulfill the diagnostic criteria

* Reproduced with permission.²³

insufficiency (0.01%) in 6947 urologic surgical procedures. Five cases (0.1%) of adrenal insufficiency were found in 4346 cardiac surgical procedures.⁵⁴ In a series of 62,473 anesthetic administrations, glucocorticoid coverage was required in 419 cases (0.7%).⁵⁵ Only 3 episodes (0.7%) of hypotension due to inadequate glucocorticoid coverage were noted.

Kehlet has reviewed²³ the reported instances of death and hypotension (Table 1) attributed to perioperative adrenal insufficiency in glucocorticoid-treated patients.

Although the majority of the early cases were inconclusive and lacked biochemical confirmation of adrenal insufficiency, Kehlet's data support the concept that the problem is real and *does* occur. In fact, a number of glucocorticoid-treated patients have undergone major surgery without perioperative glucocorticoid coverage or their usual baseline therapeutic dose of glucocorticoid.^{33,54,56,57} Most of these patients had uneventful surgical courses. The likely explanation for this paradox is that most of these patients had normal preoperative bio-

chemical indices of HPA axis function, suggesting the unreliability of the historical assessment of glucocorticoid administration. Some of the patients from these series had documented HPA axis dysfunction (i.e., an inadequate plasma cortisol response to insulin-induced hypoglycemia or ACTH administration). Several of these patients developed perioperative hypotension. The small sample size and the limited longitudinal data provided prevent further analysis.^{33,56,57} Our combined clinical experience and research findings, however, lead us to conclude that 1) clinicians need to replace glucocorticoids only in an amount equivalent to the normal physiologic response to surgical stress,^{22,24,25} and 2) the risk of anesthetizing and operating on unsupplemented glucocorticoid-treated patients (prolonged use preoperatively, but none given intraoperatively) is dependent on the duration and severity of the surgery. The HPA axis response to minor surgery (defined arbitrarily as operations performed under local anesthesia and of less than 1 hour duration) is so minimal^{22,36,40} that replacement therapy beyond the patient's normal glucocorticoid dosage seems unnecessary and potentially detrimental.

In support of these conclusions, one of us (J.B. and his co-workers) prospectively studied 40 renal allograft recipients (receiving 5–10 mg prednisone/day, long term) who required readmission to the hospital for operation or acute illness.⁵⁸ Blood samples for serum cortisol and ACTH, and “24-hour urine” collections for free cortisol determinations were obtained during the first 36 hours of hospitalization. An ACTH stimulation test was performed in 38 of the 40 patients at a later time, before discharge, when the patients were not acutely ill. The ACTH stimulation tests were abnormal in 63% of the patients. Ninety-seven per cent of patients, however, had normal or increased urinary cortisol concentrations, suggesting that circulating cortisol concentrations were sufficient to meet the patients' requirements during the time of stress.⁵⁸ There were 12 episodes of hypotension or hyponatremia. All patients responded promptly to treatment other than glucocorticoid administration. Each episode was explained readily by processes other than adrenal insufficiency. Throughout the hospitalization, none of these 40 patients received more than their baseline glucocorticoid therapy (5–10 mg prednisone/day).

These data suggest that baseline glucocorticoid therapy is sufficient to prevent adrenal insufficiency in the perioperative or stressful period. Similar to patients receiving prolonged glucocorticoid therapy, a large percentage of renal allograft recipients have biochemical evidence (i.e., an ACTH stimulation test) of inadequate adrenal reserve; nonetheless, supraphysiologic (stress) doses of glucocorticoids appear to be unnecessary. The

point needs confirmation by larger studies. The preliminary data, however, are persuasive.

ASSESSMENT OF HPA FUNCTION

The degree of HPA dysfunction depends on the amount and duration of prior glucocorticoid therapy. Medication histories, however, are notoriously inaccurate.^{23,33,36,60–63} Thus, assessment of the functional states of the HPA axis must be based on laboratory determinations. Because of the episodic nature of human cortisol secretion,^{52,63} its circadian rhythm,⁴⁸ and the variability of the HPA axis response to stress,^{23,64} single measurements of plasma cortisol, unless they are high, inadequately define the state of the HPA axis. A single plasma cortisol concentration of greater than 500 nmol/L is a strong argument for adequate HPA axis function. Most random plasma cortisol concentrations in normal subjects, however, are less than this value. Thus, to ensure that the plasma cortisol value reflects adrenal competence, a maximally stimulated value is essential.

The available provocative tests measure the plasma cortisol response to the administration of ACTH, corticotropin releasing hormone (CRH), pyrogen, lysine vasopressin, metyrapone, and insulin-induced hypoglycemia. Kehlet²³ states that the insulin tolerance and pyrogen tests are the “true stress” tests because their sites of stimulation are central. The intravenous administration of regular insulin (0.10–0.15 units/kg) results in a rapid (10–20 minutes to nadir in glucose) lowering of blood glucose concentration (usually to <2.2 mmol/L). The insulin-induced hypoglycemia triggers the release of ACTH from the pituitary gland⁶⁵ and, subsequently, cortisol from the adrenal cortex.⁶⁶ Several investigators have provided clinical evidence^{33,67,68} of the similarity of plasma cortisol responses to insulin-induced hypoglycemia and to surgery. In a review of 6580 insulin tolerance tests⁵⁹ Fish and colleagues found only six cases with untoward side effects (two episodes each of coma, angina, and impending seizure). All were reversed with intravenous glucose administration. The insulin tolerance test must be performed in a standard, closely monitored setting.

Kehlet and colleagues^{23,34,36,38,57,69,70} have provided convincing evidence that the 30-minute ACTH test is the most convenient and accurate diagnostic tool for preoperative evaluation of HPA axis function. Synthetic ACTH (Cortrosyn, Cosyntropin), in a dose of 250 µg, is administered intravenously, and a blood sample for plasma cortisol is collected 30 minutes later. Traditionally, a plasma cortisol concentration of >500 nmol/L (18–20 µg/dl) defines adequate adrenal function.

The ACTH stimulation test has a close correlation with the insulin tolerance test ($r = 0.92$; $p < 0.0001$).⁶⁹

Some authors have noted discordant results between the two tests in patients with acute pituitary disorders.^{71,72} Lindholm and Kehlet studied 200 patients with suspected hypothalamic-pituitary disorders and found a correlation between the two tests ($r = 0.83$; $p < 0.0001$).⁷³ They again confirmed the close correlation between the insulin tolerance and ACTH stimulation tests. They noted that discordant results can be obtained in an initial 1- to 2-week period after acute ACTH deprivation (pituitary surgery), a finding that can explain the normal cortisol response to ACTH in patients with acute pituitary disorders.^{71,72} We advocate use of the ACTH stimulation test as a preoperative screening test for evaluation of HPA integrity.

PRESCRIBING PERIOPERATIVE GLUCOCORTICOID COVERAGE

In a review of glucocorticoid therapy, Axelrod states that anyone who has "received a glucocorticoid in doses equivalent to 20 to 30 mg of prednisone per day for more than a week should be suspected of having HPA suppression."⁷⁴ Thus, the need to evaluate the HPA axis is a frequent consideration. Determining the actual need for glucocorticoid coverage is desirable and appropriate. The clinical complications of glucocorticoid therapy are reviewed elsewhere.⁷⁵ Special concerns about glucocorticoid use in surgical patients include the potential for adverse effects on wound healing,⁷⁶ immune function,⁷⁷⁻⁷⁹ and interaction with nondepolarizing, neuromuscular antagonists.⁸⁰⁻⁸¹

If biochemical evaluation demonstrates inadequate HPA axis function (i.e., an abnormal ACTH stimulation test), or if no preoperative testing is performed, perioperative glucocorticoid coverage should be provided. In urgent or emergent situations, the physician usually must rely on the physical signs of Cushing's syndrome that correlate with HPA axis suppression in glucocorticoid-treated patients and nonspecific laboratory information suggestive of adrenal dysfunction, such as hyponatremia and eosinophilia (indicative of the possibility of adrenal insufficiency).

The question remains — how much glucocorticoid, how often, for how long, and via what route of administration? A clue to the appropriate glucocorticoid replacement dosage is provided by the normal cortisol secretion rate in response to surgery. Kehlet²³ estimates that adults secrete 75–150 mg a day in response to major surgery and 50 mg a day during minor procedures. Table 2 reviews the available data on cortisol secretion rates and major surgery. We conclude from these data that cortisol secretion in the first 24 hours after surgery rarely exceeds 200 mg. Additionally, the secretion rate seems to parallel the duration and extent of surgery. There are

no data supporting the concept that exceeding this amount of glucocorticoid administration is ever beneficial.

Prior Therapeutic Strategies

Kehlet's⁸² recommendations for perioperative glucocorticoid coverage are based on the normal cortisol response to surgical stress. In long-term glucocorticoid-treated patients scheduled for major surgery, Kehlet⁸² recommends 25 mg of intravenous cortisol, or its equivalent, with induction of anesthesia and 100 mg given by continuous infusion over the next 24 hours. An alternative is 25 mg of cortisol equivalent given as an intravenous bolus (over 10–20 minutes) every 4 hours. After minor surgery, he recommends that the usual oral glucocorticoid replacement dose be reinstated immediately after the operation.

Lloyd⁸³ suggests minimal coverage with supplementation only if postoperative hypotension occurs. Plumpton and co-workers⁸⁴ showed that 100 mg of intramuscular hydrocortisone hemisuccinate administered every 6 hours resulted in plasma cortisol concentrations similar to those values in normal subjects undergoing similar operative stress. Gran and Pahle⁸⁵ administered a single intramuscular injection of depot betamethasone (Celestone chronodose) to glucocorticoid-treated patients. In their study, patients receiving glucocorticoids up to the time of surgery were given 2 mL of the depot agent (equivalent to 400 mg cortisol) for major surgery and 1 mL for minor surgery.⁸⁵ The agent had effects lasting up to 6 days from one injection.⁸⁴

Postoperatively, Kehlet^{23,82} recommends an intravenous infusion of cortisol equivalent (100 mg/24 hours dissolved in 500–1000 mL of isotonic saline), using return of gastrointestinal function as the indicator for return to the preoperative regimen. Although it has been customary to reduce the quantity of glucocorticoid coverage gradually over a period of days during the postoperative period, there is little evidence to support the practice, except in cases of high-dose glucocorticoid administration for prolonged periods. If HPA axis suppression is suspected postoperatively, the ACTH stimulation test will establish the diagnosis.^{53,89}

NEW RECOMMENDATIONS FOR GLUCOCORTICOID COVERAGE

We believe that glucocorticoid administration during stress should be based on the magnitude of the stress and the known glucocorticoid production rate associated with it. For **minor surgical stress** (i.e., inguinal herniorrhaphy) the daily cortisol secretion rate and static plasma cortisol measurements²⁴⁻²⁵ suggest that the glucocorti-

**Table 2. CORTISOL SECRETION RATE FOLLOWING MAJOR SURGERY—
THE OBSERVATIONS OF VARIOUS INVESTIGATORS**

Authors	Year	Cortisol Secretion Rate	Comments
Hardy and Turner ³²	1957	111 mg/24 hrs	Done by adrenal vein catheterization Constant secretion rate assumed Venous outlets overlooked 10 patients studied
Hume et al. ³⁹	1962	95 mg/24 hrs	Done by adrenal vein catheterization Constant secretion rate assumed Venous outlets overlooked 7 patients studied with indwelling adrenal vein catheters
Wise et al. ¹⁰⁵	1975	60 mg/24 hrs	Estimated from repeated plasma cortisol determinations Assumes a constant distribution volume for cortisol 3 patients studied
Kehlet and Binder ⁵⁷	1973	10 mg/hour immediately postoperative 5 mg/hr 24 hrs later 75–150 mg/24 hrs	Used ³ H cortisol before and then 2, 10, and 24 hrs after skin incision 4 patients studied
Peterson et al. ^{106,107}	1959–1960	167 mg/24 hrs Range 116–200 mg/24 hrs	Examined plasma clearance in 5 normal subjects
Ichikawa ¹⁰⁸	1966	93 mg/24 hrs 156 mg/24 hrs	Measured cortisol secretion in 2 normal subjects
Thomas et al. ¹⁰⁹	1971	222 mg/24 hrs Range 135–310 mg/24 hrs	9 asthmatic patients in which maximal cortisol secretion was studied Assumed steady state. Overestimate likely

coid target is about **25 mg of hydrocortisone equivalent**. For example, a patient receiving 5 mg of prednisone every other day for asthma who is scheduled for an inguinal hernia repair should receive preoperatively 5 mg of prednisone, with both the surgeon and anesthesiologist aware of the possible complications. If the postoperative course is uncomplicated, the patient can be returned the next day to his/her usual glucocorticoid dosage.

For **moderate surgical stress**, (i.e., nonlaparoscopic cholecystectomy, lower extremity revascularization, segmental colon resection, total joint replacement, abdominal hysterectomy), cortisol production rates suggest the glucocorticoid target is about **50 to 75 mg per day of hydrocortisone equivalent for 1 to 2 days**. For example, a patient receiving 10 mg of prednisone for systemic lupus erythematosus should receive 10 mg of prednisone (or parenteral equivalent) preoperatively and 50 mg of hydrocortisone administered intravenously intraoperatively. We suggest that 60 mg of hydrocortisone (20 mg every 8 hours) be administered intravenously on the first postoperative day and that patients return to their preoperative glucocorticoid doses (enterally or parenterally) on postoperative day 2.

Finally, for **major surgical stress** (pancreatoduodenectomy, esophagogastrectomy, total proctocolectomy, cardiac surgery involving cardiopulmonary bypass) the glucocorticoid target should be **100 to 150 mg of hydro-**

cortisone equivalent per day for 2 to 3 days. A patient with ulcerative colitis who has been receiving 40 mg of prednisone daily for several years who is scheduled for major surgery (e.g., total proctocolectomy), should receive 40 mg of prednisone (or the parenteral equivalent) preoperatively (within 2 hours of surgery), and 50 mg hydrocortisone intravenously every 8 hours after the initial dose for the first 48 to 72 hours after surgery. As a second example, a patient with arthritis who has been receiving 5 mg of prednisone daily and who is undergoing similar major surgery should require 5 mg oral prednisone (or the hydrocortisone equivalent intravenously) as a preoperative dose, with 25 mg of hydrocortisone to be given intraoperatively, and 25 mg to be administered in the 8 hours after surgery. Over the subsequent 48 hours, we would prescribe 25 mg of hydrocortisone every 8 hours. The examples provided for major surgical coverage demonstrate our belief that the provision of perioperative glucocorticoid coverage must account for the patient's preoperative glucocorticoid dose and the duration and severity of surgery or other stress.

To our knowledge, there are no data that indicate these recommended equivalent doses need to be exceeded. Thus, the patient who is receiving a maintenance dose of glucocorticoid therapy that exceeds the estimated stress requirement will not need more glucocorticoid coverage during the stress period. After uncomplicated major sur-

gery, plasma cortisol concentrations decrease rapidly. The circulating cortisol concentration is normal by 24 to 48 hours after surgical stress in most patients.^{22,87,88} A postoperative increase in cortisol secretion is presumptive evidence for a continued or new stressor (i.e., fever, peritonitis, etc.). In the case of postoperative complications, we recommend continued glucocorticoid administration consistent with the postoperative stress response.

THE FUTURE

Although the HPA response to stress is well defined and Addisonian patients or adrenalectomized animals succumb to stress, the precise physiologic roles of glucocorticoids in meeting the demands of stress are unknown. We believe that recent work⁹⁰⁻¹⁰⁴ has begun to identify the mechanisms by which glucocorticoids exert their actions during stress. Some of this work has focused on the relationship of and communication between the immune system (lymphocyte and macrophage-borne cytokines) and the HPA axis.

Woloski and colleagues⁹⁰ provided experimental evidence that monokines, such as interleukin-1 (IL-1), when added to cultures of mouse pituitary tumor cells and adrenal cells, stimulate the release of ACTH and glucocorticoids. They suggest that increases in IL-1 during states of inflammatory/immune response (stress) may be responsible for activation of the HPA axis, and hypothesized that a feedback loop exists between the immune system (monokines) and the HPA axis. Other investigators have noted IL-1-induced secretion of ACTH by rat pituitary cells.⁸⁹⁻⁹⁰ Besedovsky and co-workers⁹³ showed that infusions of IL-1 in mice were associated with increased blood levels of ACTH and glucocorticoids. Further, these investigators and others⁹³⁻⁹⁶ suggest that corticotropin-releasing factor modulates the IL-1-induced pituitary adrenal activation. Recent experimental evidence suggests that interleukin-6 is a potent stimulus for release of ACTH *in vitro*⁹⁷ and *in vivo*.⁹⁸

Infusions of tumor necrosis factor (TNF) cause sustained increases in circulating glucocorticoid concentrations in rats⁹⁹ and dogs.¹⁰⁰ Adrenalectomized rats have a 100% mortality rate from such an infusion.⁹⁹ Pre-treating these animals with dexamethasone (0.3 mg/kg) prevents death from TNF or IL-1 infusions¹⁰¹ and inhibits endotoxin-induced TNF release.¹⁰²

Marano and colleagues¹⁰³ measured TNF and serum cortisol concentrations in patients with severe burn injury. In 16 patients studied longitudinally, they found that cortisol levels are significantly higher ($p < 0.02$) in the absence of detectable circulating levels of TNF.¹⁰³ They suggest that high cortisol levels may interfere with cytokine production.¹⁰³ Naito and co-workers¹⁰⁴ longitudinally

evaluated circulating ACTH, cytokines (TNF, IL-6), and cortisol concentrations in patients undergoing pancreatoduodenectomy or total joint replacement procedures. They showed that (in pancreatoduodenectomy patients) significant intraoperative and postoperative increases in circulating cytokine concentrations were related temporally to elevations in postoperative cortisol values in the first 48 to 72 hours after surgery.¹⁰⁴ These studies suggest a connection between the stress response and the pro-inflammatory cytokines. Further studies should define the effect of glucocorticoids (during surgery/stress) as counterregulatory hormones modulating the release or modifying the systemic actions of cytokines.

Four decades have passed since a standard of care was established for perioperative glucocorticoid coverage. This initial standard was based on anecdotal information. Experimental data accumulated since that time support the concept that new recommendations are in order. This article provides a framework for those new recommendations in the context of the need to individualize this important aspect of pharmacologic management.

References

1. Fraser CG, Preuss FS, Bigford WD. Adrenal atrophy and irreversible shock associated with cortisone therapy. *JAMA* 1952; 149: 1542-1543.
2. Addison T. On the constitutional and local effects of diseases of the suprarenal capsules. London: Samuel Highley, 1855.
3. Cushing H. The pituitary body and its disorders. Philadelphia: JB Lippincott, 1912.
4. Cushing H. The basophil adenomas of the pituitary body and their clinical manifestations (pituitary basophilism). *Bull Johns Hopkins Hosp* 1932; 50:137.
5. Swingle WW, Pfiffner JJ, Vars HM, et al. The function of the adrenal cortex hormone and the cause of death from adrenal insufficiency. *Science* 1933; 77:58-64.
6. Swingle WW, Parkins WM, Taylor AR, Hays HW. A study of the circulatory failure and shock following trauma to the healthy, vigorous adrenalectomized dog. *Am J Physiol* 1938; 124:22-29.
7. Swingle WW, Hays HW, Remington JW, et al. The effect of primary doses of desoxycorticosterone acetate in preventing circulatory failure and shock in the adrenalectomized dog. *Am J Physiol* 1949; 132:249-258.
8. Swingle WW, Remington JW, Drill VS, Kleinberg W. Differences among adrenal steroids with respect to their efficacy in protecting the adrenalectomized dog against circulatory failure. *Am J Physiol* 1942; 136:567-576.
9. Swingle WW, DaVanzo JP, Crossfield HC, et al. Glucocorticoids and maintenance of blood pressure and plasma volume of adrenalectomized dogs subjected to stress. *Proc Soc Exp Biol Med* 1959; 100:617-622.
10. Langley JN. Observations on the physiologic action of extracts of the suprarenal bodies. *J Physiol (London)* 1901; 27:237-256.
11. Lewandowsky M. Ueber eine Wirkung des Nebennieren extractes auf das Auge. *Zentralb Physiol* 1898; 12:599-600.

12. Swingle WW, Pfiffner JJ. Experiments with an active extract of the suprarenal cortex. *Anat Rec* 1930; 44:225–226.
13. Swingle WW, Pfiffner JJ. An aqueous extract of the suprarenal cortex which maintains the life of bilaterally adrenalectomized cats. *Science* 1930; 71:321–322.
14. Reichstein T, Shoppee CW. The hormones of the adrenal cortex. In Harris RS, Thimann KV. *Vitamins and Hormones: Advances in Research and Applications*. New York: Academic Press, 1943, 1:346–413.
15. De Fremery P, Laquer E, Reichstein T, et al. Corticososteron, a crystalloid compound with biological activity of the adrenal-cortical hormone. *Nature* 1937; 139:26.
16. Kendall EC, Mason HL, Hoehn WM, McKenzie BF. The structure and physiologic activity of Compound B: its relation to Compound A and Reichsteins cortisterone. *Proc Staff Meet Mayo Clin* 1937; 12:136–140.
17. Mason HL, Hoehn WM, Kendall EC. Chemical studies of the suprenal cortex, structures of compounds C, D, E, F, and G. *J Biol Chem* 1938; 124:459–474.
18. Hench PS, Kendall EC, Slocumb CH, Polley HF. The effect of a hormone of the adrenal cortex (17-hydroxy-11-dehydrocorticosterone; compound E) and of pituitary adrenocorticotrophic hormone on rheumatoid arthritis. *Proc Staff Meet Mayo Clin* 1949; 24:181–197.
19. Hench PS, Slocumb CH, Polley HF, Kendall EC. Effect of cortisone and pituitary adrenocorticotrophic hormone (ACTH) on rheumatic diseases. *JAMA* 1950; 144:1327–1335.
20. Lewis L, Robinson RF, Yee J, et al. Fatal adrenal cortical insufficiency precipitated by surgery during prolonged continuous cortisone infusion. *Ann Intern Med* 1953; 39:116–125.
21. Esteban N, Loughlin T, Yergey A, et al. Daily cortisol production rate in man determined by stable isotope dilution, mass spectrometry. *J Clin Endocrinol Metab* 1991; 71:39–45.
22. Chernow B, Alexander HR, Thompson WR, et al. The hormonal responses to surgical stress. *Arch Intern Med* 1987; 147:1273–1278.
23. Kehlet H. *Clinical Course and Hypothalamic-Pituitary-Adrenocortical Function in Glucocorticoid-Treated Surgical Patients*. Copenhagen: FADL's Forlag, 1976.
24. Udelsman R, Ramp J, Gallucci WT, et al. Adaptation during surgical stress—a re-evaluation of the role of glucocorticoids. *J Clin Invest* 1986; 44:1377–1381.
25. Udelsman R, Goldstein DS, Loriaux DL, Chrousos GP. Catecholamine-glucocorticoid interactions during surgical stress. *J Surg Res* 1987; 43:539–545.
26. Chernow B, Anderson DM. Endocrine responses to critical illness. *Semin Resp Med* 1985; 7:1–10.
27. Meguid MM, Brennan MF, Aoki TT, et al. Hormone-substrate interrelationship following trauma. *Arch Surg* 1974; 109:776–783.
28. Carey LC, Clontier CT, Lowery BD. Growth hormone and adrenal cortical response to shock and trauma in the human. *Ann Surg* 1971; 174:451.
29. Carey LC, Lowery BD, Clontier CT. Blood sugar and insulin response in shock. *Ann Surg* 1970; 172:342.
30. Cooper CE, Nelson DH. ACTH levels in plasma in preoperative and surgically stressed patients. *J Clin Invest* 1962; 41:1599–1605.
31. Yalow RS, Varrano-Aharon N, Echemendia E, Berson S. HCG and ACTH secretory response to stress. *Horm Metab Res* 1969; 1:3–8.
32. Hardy JD, Turner MD. Hydrocortisone secretion in man: studies of adrenal vein blood. *Surgery* 1957; 42:194–201.
33. Plumpton FS, Besser GM, Cole PV. Corticosteroid treatment and surgery—An investigation of the indications for steroid cover. *Anesthesia* 1969; 24:3–11.
34. Kehlet H, Binder C, Engbaek C. Imitation of the adrenocortical response to surgery by intravenous infusion of synthetic human ACTH. *Acta Endocrinologica (Copenhagen)* 1973; 72:75–80.
35. Banks P. The adreno-cortical response to oral surgery. *Br J Oral Surg*; 1970; 8:32–44.
36. Kehlet H, Binder C. Adrenocortical function and clinical course during and after surgery in unsupplemented glucocorticoid-treated patients. *Br J Anaesth* 1973; 45:1043–1048.
37. Dorner G, Stahl F, Wendt F, Schädlich M. Beurteilung der Nebennierenrinden funktion beim Operations stress durch fluorometrische Bestimmung von unkonjugiertem Lortisol und Corticosteron in Plasma und Harn. *Brun's Bietrage zur Klinischen Chirurgie* 1966; 212:467–480.
38. Espiner EA. Urinary cortisol excretion in stress situations and in patients with Cushing's syndrome. *J Endocrinol* 1966; 35:29–44.
39. Hume DM, Bell CC, Bartter F. Direct measurement of adrenal secretion during operative trauma and convalescence. *Surgery* 1962; 52:174–187.
40. Hume DM, Nelson DH, Miller DW. Blood and urinary 17-hydroxycorticosteroids in patients with severe burns. *Ann Surg* 1956; 143:316–329.
41. Kleg HK, Peerenboom H, Strohmeyer G, Druskemper HL. Cortisol excretion into gastric juice—studies in health, in digestive ulcer disease and in surgery stress. *Dig Dis Sci* 1983; 28:494–501.
42. Lacoumenta S, Paterson JL, Burrin J, et al. Effects of two differing halothane concentrations on the metabolic and endocrine responses to surgery. *Br J Anaesth* 1986; 58:844–850.
43. Lehtinen AM, Fyhrquist F, Kivalo I. The effect of fentanyl on arginine vasopressin and cortisol secretion during anesthesia. *Anesth Analg* 1984; 63:25–30.
44. Beyer HS, Parker L, Li CH, et al. β -endorphin attenuates the serum cortisol response to exogenous adrenocorticotropin. *J Clin Endocrinol Metab* 1986; 62:808–811.
45. Downing R, Davis I, Black J, Windson CWO. Effect of intrathecal morphine on the adrenocortical and hyperglycemic responses to upper abdominal surgery. *Br J Anaesth* 1986; 58:858–861.
46. Engquist S, Brandt MR, Fernandes A, Kehlet H. The blocking effect of epidural analgesia on the adrenocortical and hyperglycemic response to surgery. *Acta Anaesth Scand* 1977; 21:330–335.
47. Rittmaster RS, Cutler GB Jr, Sobel DO, et al. Morphine inhibits the pituitary-adrenal response to ovine corticotropin-releasing hormone in normal subjects. *J Clin Endocrinol Metab* 1985; 60:891–895.
48. Masala A, Satta G, Alagna S, et al. Effect of clonidine on stress-induced cortisol release in man during surgery. *Pharmacol Res Commun* 1985; 17:293–298.
49. Sherman B, Wysham C, Pfohl B. Age-related changes in the circadian rhythm of plasma cortisol in man. *J Clin Endocrinol Metab* 1985; 61:439–443.
50. Weitzman ED, Zimmerman JL, Czesler CA, Ronda J. Cortisol secretion is inhibited during sleep in normal man. *J Clin Endocrinol Metab* 1983; 56:352–358.
51. Catalano RD, Parameswaran V, Ramachandran J, Trunkey DD. Mechanisms of adrenocortical depression during *Escherichia Coli* shock. *Arch Surg* 1984; 119:145–150.
52. Avgerinos PC, Schurmeyer TH, Gold PW, et al. Pulsatile administration of human corticotropin-releasing hormone in patients with secondary adrenal insufficiency—restoration of the normal cortisol secretory pattern. *J Clin Endocrinol Metab* 1986; 62:816–821.
53. Mohler JL, Michael KA, Freedman AM, et al. The evaluation of

- postoperative function of the adrenal gland. *Surg Gynecol Obstet* 1985; 161:551-556.
54. Alford WC Jr, Meador CK, Mihalevich J, et al. Acute adrenal insufficiency following cardiac surgical procedures. *J Thorac Cardiovasc Surg* 1979; 78:489-493.
55. Kaalund Jensen J, Elb S. Preoperative and postoperative complications in previously corticosteroid-treated patients. *Nordisk Medicin* 1966; 25(8):978.
56. Knudsen L, Christiansen LA, Lorentzen JE. Hypotension during and after operation in glucocorticoid-treated patients. *Br J Anaesth* 1981; 53:295-301.
57. Kehlet K, Binder C. Value of an ACTH test in assessing hypothalamic-pituitary-adrenocortical function in glucocorticoid-treated patients. *Br Med J* 1973; 2:147-149.
58. Bromberg JS, Alfrey EJ, Barker CF, et al. Adrenal suppression and steroid supplementation in renal transplant recipients. *J Transplantation* 1991; 51:385-390.
59. Fish HR, Chernow B, O'Brian JT. Endocrine and neurophysiologic responses of the pituitary to insulin-induced hypoglycemia: A review. *Metabolism* 1986; 35:763-780.
60. Marks LJ, Donovan MJ, Duncan FJ, Karger R. Adrenocortical response to surgical operation in patients treated with corticosteroids or corticotropin prior to surgery. *J Clin Endocrinol Metab* 1959; 19:1458-1470.
61. Tredwell BLJ, Savage O, Sever ED, Copeman WSC. Pituitary adrenal function during corticosteroid therapy. *Lancet* 1963; i: 355-358.
62. Daly JR, Glass D. Corticosteroid and growth hormone response to hypoglycemia in patients on long term treatment with corticotropin. *Lancet* 1971; i:476-477.
63. Schlaghecke R, Kormely E, Santen RT, Rudderskamp P. The effect of long-term glucocorticoid therapy on pituitary-adrenal responses to exogenous corticotropin-releasing hormone. *N Engl J Med* 1992; 326:226-230.
64. Reincke M, Allolio B, Wurth G, Winkelmann W. The Hypothalamic-Pituitary-Adrenal Axis in critical illness: response to dexamethasone and corticotropin-releasing hormone. *J Clin Endocrinol Metab* 1993; 77:151-156.
65. Berson SA, Yalow RS. Radioimmunoassay of ACTH in plasma. *J Clin Invest* 1968; 47:2725-2751.
66. Landon J, Wynn V, James VHT. Adrenocortical response to insulin-induced hypoglycemia. *J Endocrinol* 1963; 27:183-192.
67. Plumpton FS, Besser GM. The adrenocortical response to surgery and insulin-induced hypoglycemia in corticosteroid-treated and normal subjects. *Br J Surg* 1969; 56:216-219.
68. Blichert-Toft M, Christianson C, Enhquist A, et al. Comparison of pituitary-adrenocortical response to hypoglycemia and surgery. *Acta Anaesth Scand* 1979; 23:103-106.
69. Lindholm J, Kehlet H, Blichert-Toft M, et al. Reliability of the 30-minute ACTH test in assessing hypothalamic-pituitary-adrenal function. *J Clin Endocrinol Metab* 1978; 47:272-274.
70. Hjortrup A, Kehlet J, Lindholm L, Stentoft P. Value of the 30-minute ACTH test in demonstrating hypothalamic-pituitary-adrenocortical insufficiency after acute ACTH deprivation. *J Clin Endocrinol Metab* 1983; 57:668-670.
71. Borst G, Michienfelder H, O'Brian J. Discordant cortisol response to exogenous ACTH and insulin-induced hypoglycemia in patients with pituitary disease. *N Engl J Med* 1982; 306:1462-1464.
72. Reschini E, Catania A, Giustina G. Plasma Cortisol response to ACTH does not accurately indicate the state of hypothalamic-pituitary-adrenal axis. *J Endocrinol Invest* 1982; 5:259-261.
73. Lindholm L, Kehlet H. Re-evaluation of the clinical value of the 30-minute ACTH test in assessing the hypothalamic-pituitary-adrenocortical function. *Clin Endocrinol* 1987; 26:53-59.
74. Axelrod L. Glucocorticoid therapy. *Medicine* 1976; 55:39-63.
75. Dujovne CA, Azarnoff DL. Clinical complications of corticosteroid therapy—a review. *Med Clin N Amer* 1973; 57:1331-1342.
76. Goforth P, Gudas CJ. Effects of steroids on wound-healing—a review of the literature. *J Foot Surg* 1980; 19:22-28.
77. Schaffner A. Therapeutic concentrations of glucocorticoids suppress the antimicrobial activity of human macrophages without impairing their responsiveness to gamma interferon. *J Clin Invest* 1985; 76:1755-1764.
78. Baltch AL, Hammer MC, Smith RP, et al. Comparison of the effect of three adrenal corticosteroids on human granulocyte function against *Pseudomonas aeruginosa*. *J Trauma* 1986; 26: 525-533.
79. Goodwin JS, Atluru D, Dierakowski S, Lianos EA. Mechanism of action of glucocorticoids—inhibition of T-cell proliferation and interleukin-2 production by hydrocortisone is reversed by leukotriene B₄. *J Clin Invest* 1986; 77:1244-1250.
80. Laffin MJ. Interaction of pancuronium and corticosteroids. *Anesthesiology* 1977; 47:471-472.
81. Azar I, Kumar D, Betcher AM. Resistance to pancuronium in an asthmatic patient treated with aminophylline and steroids. *Can Anaesth Soc J* 1982; 29:280-282.
82. Kehlet H. A rational approach to dosage and preparation of parenteral glucocorticoid substitution therapy during surgical procedures. *Acta Anaesth Scand* 1975; 19:260-264.
83. Lloyd EF. A rational regimen for perioperative steroid supplements and a clinical assessment of the requirement. *Ann Royal Coll Surg Engl* 1981; 63:54-57.
84. Plumpton FS, Besser GM, Cole PV. Corticosteroid treatment and surgery. *Anaesthesia* 1969; 24:12-18.
85. Gran L, Pahle JA. Rational substitution for steroid treated patients. *Anaesthesia* 1978; 33:59-61.
86. Frey H, Norman N. Duration of action of depot corticosteroids. *Pharmacologia Clinica* 1970; 2:109-112.
87. Jasani M, Freeman P, Boyle J, et al. Studies of the rise in plasma 11-hydroxycorticosteroids (11-OHCS) in corticosteroid-treated patients with rheumatoid arthritis during surgery: correlations with the functional integrity of the hypothalamo-pituitary-adrenal axis. *Quart J Med* 1968; 37:407-421.
88. Mattingly D, Tyler C. Plasma 11-hydroxycorticoid levels in surgical stress. *Proc Royal Soc Med* 1965; 58:24-26.
89. Beyer HS, Bantle JP, Mariash CN, et al. Use of the dexamethasone-adrenocorticotropin test to assess the requirement for continued glucocorticoid replacement therapy after pituitary surgery. *J Clin Endocrinol Metab* 1985; 60:1012-1018.
90. Woloski BM, Smith EM, Meyey WJ, et al. Corticotropin-releasing activity of monokines. *Science* 1985; 230:1037-1037.
91. Bernton EW, Beach JE, Holaday JW, et al. Release of multiple hormones by a direct action of interleukin-1 on pituitary cells. *Science* 1987; 238:519-521.
92. Beach JE, Smallridge RC, Kinzer CA, et al. Rapid release of multiple hormones from rat pituitaries perfused with recombinant interleukin-1. *Life Sci* 1989; 44:1-7.
93. Besedovsky H, Del Rey A, Sorkin E, Dinarello CA. Immunoregulatory feedback between interleukin-1 and glucocorticoid hormones. *Science* 1986; 233:652-654.
94. Berkenbosch F, Van Oers J, Del Rey A, et al. Corticotropin releasing factor-producing neurons in the rat activated by interleukin-1. *Science* 1987; 238:524-526.
95. Uehara A, Gottschall PE, Dahl RR, Arimura A. Interleukin-1 stimulates ACTH release by an indirect action which requires en-

- dogenous corticotropin-releasing factor. *Endocrinology* 1987; 121:1580–1582.
96. Breder CD, Dinarello CA, Saper CB. Interleukin-1 immunoreactive innervation of the human hypothalamus. *Science* 1988; 240: 321–323.
97. Spangelo BL, MacLeod RM, Isakson PC. Production of interleukin-6 by anterior pituitary cells in vitro. *Endocrinology* 1990; 126:582–586.
98. Naitoh Y, Fukata J, Tominaga T, et al. Interleukin-6 stimulates the secretion of adrenocorticotrophic hormone in conscious, freely-moving rats. *Biochem Biophys Res Comm* 1988; 155:1459–1463.
99. Mealy KM, van Lanschot JJB, Robinson BG, et al. Are the catabolic effects of tumor necrosis factor mediated by glucocorticoids? *Arch Surg* 1990; 125:42–48.
100. Tracey KJ, Lowry SF, Fahey TJ, et al. Cachectin/tumor necrosis factor induces lethal shock and stress hormone responses in the dog. *Surg Gynecol Obstet* 1987; 164:415–422.
101. Bertini R, Bianchi M, Ghezzi P. Adrenalectomy sensitizes mice to the lethal effects of interleukin-1 and tumor necrosis factor. *J Exp Med* 1988; 167:1708–1712.
102. Malcolm D, Hinson D, Chia J. Dexamethasone inhibits endotoxin-induced tumor necrosis factor release in vivo. *Circ Shock* 1988; 24:296 (Abstract).
103. Marano MA, Fong Y, Moldawer LL, et al. Serum cachectin/tumor necrosis factor in critically ill patients with burns correlates with infection and mortality. *Surg Gynecol Obstet* 1990; 170:32–38.
104. Naito Y, Tamai S, Shingu K, et al. Responses of plasma adrenocorticotrophic hormone, cortisol, and cytokines during and after upper abdominal surgery. *Anesthesiology* 1992; 77:426–431.
105. Wise L, Margraf HW, Ballinger WF. A new concept on the pre- and postoperative regulation of cortisol secretion. *Surgery* 1972; 72:290–299.
106. Peterson RE. The miscible pool and turnover rate of adrenocortical steroids in man. *Recent Prog Hormone Res* 1959; 15:231–274.
107. Peterson RE, Nokes G, Chen PS, Black RL. Estrogens and adrenocortical function in man. *J Clin Endocrinol Metab* 1960; 20: 495–514.
108. Ichikawa Y. Metabolism of cortisol-4-C14 in patients with infections and collagen diseases. *Metabolism* 1966; 15:613–625.
109. Thomas JP, El-Shaboury AH. Aldosterone secretion in steroid treated patients with adrenal suppression. *Lancet* 1917; i:623–625.